

Patent claims: as submitted November 9, 2000

1. An active ingredient combination comprising tris(2-ethylhexyl) 4,4',4''-(1,3,5-triazine-2,4,6-triyltriimino)trisbenzoate and one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols.
2. Method of using one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols as solvent, solubility promoter or solubilizer for tris(2-ethylhexyl) 4,4',4''-(1,3,5-triazine-2,4,6-triyltriimino)trisbenzoate, in particular ~~for the use in~~ light protection compositions.
3. The combination as claimed in claim 1 , characterized in that the total amount of tris(2-ethylhexyl) 4,4',4''-(1,3,5-triazine-2,4,6-triyltriimino)trisbenzoate in the finished cosmetic or dermatological preparations is chosen from the range 0.1-10.0% by weight, preferably 0.5-6.0% by weight, in each case based on the total weight of the preparations.
4. The combination as claimed in claim 1 , characterized in that the ester(s) of branched-chain carboxylic acids and branched-chain alcohols is / are present in a mixture with the parent alcohols R_2-OH .
5. The combination as claimed in claim 1 , characterized in that the ester of branched-chain carboxylic acids and branched-chain alcohols chosen is hexyldecyl laurate.
6. The combination as claimed in claim 1 , characterized in that the total amount of one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols in the finished cosmetic or dermatological preparations is chosen from the range 0.1-25.0% by weight, preferably 0.5-15.0% by weight, in each case based on the total weight of the preparations.

00000103-110000

Description

Cosmetic and dermatological light protection formulations containing triazine derivatives and one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols

The harmful effect of the ultraviolet part of solar radiation on the skin is generally known. While rays with a wavelength of less than 290 nm (the so-called UVC region) are absorbed by the ozone layer in the earth's atmosphere, rays in the range between 290 nm and 320 nm, the so-called UVB range, cause erythema, simple sunburn or even burns of varying severity.

The maximum given for the erythema activity of sunlight is the narrower range around 308 nm.

Numerous compounds are known for protecting against UVB radiation; these are usually derivatives of 3-benzylidenecamphor, of 4-aminobenzoic acid, of cinnamic acid, of salicylic acid, of benzophenone and also of 2-phenylbenzimidazole.

It is also important to have available filters for the range between about 320 nm and about 400 nm, the so-called UVA region, since its rays can likewise cause damage. Thus, it has been found that UVA radiation leads to damage of the elastic and collagenous fibers of connective tissue, causing premature aging of the skin, and that it


is to be regarded as a cause of numerous phototoxic and photoallergic reactions. The harmful effect of UVB radiation can be intensified by UVA radiation.

However, UV radiation can also lead to photochemical reactions, in which case the photochemical reaction products intervene in the skin's metabolism.

Such photochemical reaction products are predominantly free-radical compounds, e.g. hydroxyl radicals. Undefined free-radical photoproducts which form in the skin itself can also display uncontrolled secondary reactions because of their high reactivity. However, singlet oxygen, a non-radical excited state of the oxygen molecule, can also arise during UV radiation, as can short-lived epoxides and many other species. Singlet oxygen, for example, differs from the normal triplet oxygen (free-radical ground state) by virtue of its increased reactivity. However, excited reactive (free-radical) triplet states of the oxygen molecule also exist.

Furthermore, UV radiation is a type of ionizing radiation. There is therefore the risk that UV exposure may also produce ionic species which then, for their part, are capable of oxidative intervention in the biochemical processes.

An advantageous UVB filter is tris(2-ethylhexyl) 4,4',4''-(1,3,5-triazine-2,4,6-triyltriimino)trisbenzoate, also known as 2,4,6-tris-[anilino(p-carbo-2'-ethyl-1'-hexyloxy)]-1,3,5-triazine.



1

The esters of unbranched-chain carboxylic acids and branched-chain alcohols according to the invention can advantageously be chosen from the group of substances of the general formula



in which R₁ is an unbranched-chain alkyl radical having 3 to 30 carbon atoms, and R₂ represents mono- or polybranched alkyl radicals having 3 to 30 carbon atoms.

R₁ is advantageously an unbranched-chain alkyl radical having up to 20 carbon atoms.

In addition, R₂ is advantageously a mono- or polybranched alkyl radical having up to 25 carbon atoms.

For the purposes of the present invention, the carboxylic ester(s) is/are very particularly advantageously present in a mixture with the parent alcohols R_2-OH .

It was nevertheless surprising that the addition of one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols effects stabilization of solutions of tris(2-ethylhexyl) 4,4',4''-(1,3,5-triazine-2,4,6-triyltriimino)trisbenzoate, since the latter substance not only has poor solubility, but also readily crystallizes out again from its solution. The invention therefore also provides a method of stabilizing solutions of tris(2-ethylhexyl) 4,4',4''-(1,3,5-triazine-2,4,6-triyltriimino)trisbenzoate, which comprises adding an effective content of one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols to such solutions.

The total amount of tris(2-ethylhexyl) 4,4',4''-(1,3,5-triazine-2,4,6-triyltriimino)trisbenzoate in the finished cosmetic or dermatological preparations is advantageously chosen from the range 0.1-10.0% by weight, preferably 0.5-6.0% by weight, based on the total weight of the preparations.

The total amount of one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols in the finished cosmetic or dermatological preparations is advantageously chosen from the range 0.1-25.0% by weight, preferably 0.5-15.0% by weight, based on the total weight of the preparations.

It is advantageous to choose weight ratios of tris(2-ethylhexyl) 4,4',4''-(1,3,5-triazine-2,4,6-triyltriimino)trisbenzoate and one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols from the range 1 : 10 to 10 : 1, preferably 1 : 4 to 4 : 1.

Cosmetic and dermatological preparations according to the invention also advantageously comprise inorganic pigments based on metal oxides and/or other

For the purposes of the present invention, it is particularly advantageous, although not imperative, if the inorganic pigments are present in hydrophobic form, i.e. they have been surface-treated to repel water. This surface treatment can consist in providing the pigments with a thin hydrophobic layer by processes known per se.

$$n \text{ TiO}_2 + m (\text{RO})_3 \text{ Si-R}' \xrightarrow{6n \text{ TiO}_2 (\text{surf.})}$$

Advantageous TiO₂ pigments are available, for example, under the trade names MT 100 T from TAYCA, also M 160 from Kemira and T 805 from Degussa.

The cosmetic and/or dermatological light protection formulations according to the invention can have the conventional compositions and can be used for cosmetic and/or dermatological light protection, and also for the treatment, care and cleansing of the skin and/or hair and as make-up products in decorative cosmetics.

For use, the cosmetic and dermatological preparations according to the invention are applied to the skin and/or hair in sufficient amount and in the manner conventional for cosmetics.

Particularly preferred cosmetic and dermatological preparations are those which are in the form of a sunscreen. Advantageously, these can additionally comprise at least one



An additional content of antioxidants is generally preferred. According to the invention, favorable antioxidants which can be used are all antioxidants suitable or customary for cosmetic and/or dermatological applications.

The antioxidants are advantageously chosen from the group consisting of amino acids (e.g. glycine, histidine, tyrosine, tryptophan) and derivatives thereof, imidazoles, (e.g. urocanic acid) and derivatives thereof, peptides, such as D,L-carnosine, D-carnosine, L-carnosine and derivatives thereof (e.g. anserine), carotenoids, carotenes (e.g. α -carotene, β -carotene, lycopene) and derivatives thereof, chlorogenic acid and derivatives thereof, lipoic acid and derivatives thereof (e.g. dihydrolipoic acid), aurothioglucose, propylthiouracil and other thiols (e.g. thioredoxin, glutathione, cysteine, cystine, cystamine and glycosyl, N-acetyl, methyl, ethyl, propyl, amyl, butyl and lauryl, palmitoyl, oleyl, γ -linoleyl, cholesteryl and glyceryl esters thereof) and salts thereof, dilauryl thiodipropionate, distearyl thiodipropionate, thiodipropionic acid and derivatives thereof (esters, ethers, peptides, lipids, nucleotides, nucleosides and salts) and sulfoximine compounds (e.g. buthionine-sulfoximines, homocysteine-sulfoximine, buthionine-sulfones, penta-, hexa-, heptathionine-sulfoximines) in very low tolerated doses (e.g. pmol to μ mol/kg), and also (metal) chelating agents (e.g. α -hydroxyfatty acids, palmitic acid, phytic acid, lactoferrin), α -hydroxy acids (e.g. citric acid, lactic acid, malic acid), humic acid, bile acid, bile extracts, bilirubin, biliverdin, EDTA, EGTA and derivatives thereof, unsaturated fatty acids and derivatives thereof (e.g. γ -linolenic acid, linoleic acid, oleic acid), folic acid and derivatives thereof, furfurylidenesorbitol and

The amount of the abovementioned antioxidants (one or more compounds) in the preparations is preferably from 0.001 to 30% by weight, particularly preferably from 0.05 to 20% by weight, in particular from 1 to 10% by weight, based on the total weight of the preparation.

If vitamin E and/or derivatives thereof are used as the antioxidant or antioxidants, their respective concentrations are advantageously chosen from the range 0.001 - 10% by weight, based on the total weight of the formulation.

If vitamin A or vitamin A derivatives or carotenes or derivatives thereof are used as the antioxidant or antioxidants, their respective concentrations are advantageously chosen from the range 0.001 - 10% by weight, based on the total weight of the formulation.

The lipid phase can advantageously be chosen from the following group of substances:

- mineral oils, mineral waxes
- oils, such as triglycerides of capric or of caprylic acid, but preferably castor oil;
- fats, waxes and other natural and synthetic fatty substances, preferably esters of fatty acids with alcohols of low carbon number, e.g. with isopropanol,

- alkyl benzoates;
- silicone oils such as dimethylpolysiloxanes, diethylpolysiloxanes, diphenylpolysiloxanes and mixtures thereof.

The oil phase can also advantageously be chosen from the group of branched and unbranched hydrocarbons and hydrocarbon waxes, of silicone oils, of dialkyl ethers, from the group of saturated or unsaturated, branched or unbranched alcohols, and also fatty acid triglycerides, namely the triglycerol esters of saturated and/or unsaturated, branched and/or unbranched alkanecarboxylic acids having a chain length of from 8 to 24, in particular 12 - 18, carbon atoms. The fatty acid triglycerides can advantageously be chosen, for example, from the group of synthetic, semi-synthetic and natural oils, e.g. olive oil, sunflower oil, soybean oil, groundnut oil, rapeseed oil, almond oil, palm oil, coconut oil, palm kernel oil and the like.

For the purposes of the present invention, any mixtures of such oil and wax components can also advantageously be used. When required, it can also be

The oil phase is advantageously chosen from the group consisting of 2-ethylhexyl isostearate, octyldodecanol, isotridecyl isononanoate, isoeicosane, 2-ethylhexyl cocoate, C₁₂-C₁₅-alkyl benzoate, caprylic/capric triglyceride and dicaprylyl ether.

For the purposes of the present invention, of the hydrocarbons, paraffin oil, squalane and squalene can advantageously be used.

Cyclomethicone (octamethylcyclotetrasiloxane) is advantageously used as silicone oil to be used according to the invention. However, other silicone oils can also be used advantageously for the purposes of the present invention, for example hexamethylcyclotrisiloxane, polydimethylsiloxane, poly(methylphenylsiloxane).

The aqueous phase of the preparations according to the invention may advantageously comprise

- alcohols, diols or polyols of low carbon number, and also ethers thereof, preferably ethanol, isopropanol, propylene glycol, glycerol, ethylene glycol, ethylene glycol monoethyl or monobutyl ether, propylene glycol monomethyl, monoethyl or monobutyl ether, diethylene glycol monomethyl or monoethyl ether and analogous products, and also alcohols of low carbon number, e.g.

ethanol, isopropanol, 1,2-propanediol, glycerol, and in particular one or more thickeners which can advantageously be chosen from the group consisting of silicon dioxide, aluminum silicates, polysaccharides and derivatives thereof, e.g. hyaluronic acid, xanthan gum, hydroxypropylmethylcellulose, and particularly advantageously from the group of polyacrylates, preferably a polyacrylate from the group of the so-called Carbopols, for example Carbopol grades 980, 981, 1382, 2984, 5984, in each case individually or in combination.

The cosmetic or dermatological light protection preparations advantageously comprise inorganic pigments, in particular micropigments, e.g. in amounts of from 0.1% by weight to 30% by weight, preferably in amounts of from 0.5% by weight to 10% by weight, but in particular from 1% by weight to 6% by weight, based on the total weight of the preparations.

It is advantageous according to the invention to use, apart from the combinations according to the invention, oil-soluble UVA filters and/or UVB filters in the lipid phase and/or water-soluble UVA filters and/or UVB filters in the aqueous phase.

The light protection formulations according to the invention can advantageously comprise further substances which absorb UV radiation in the UVB region, the total amount of filter substances being e.g. from 0.1% by weight to 30% by weight, preferably from 0.5 to 10% by weight, in particular from 1 to 6% by weight, based on the total weight of the preparations, in order to provide cosmetic preparations which protect the skin from the entire range of ultraviolet radiation. They can also be used as sunscreens.

The further UVB filters can be oil-soluble or water-soluble. Examples of advantageous oil-soluble UVB filter substances are:

- 3-benzylidenecamphor derivatives, preferably 3-(4-methylbenzylidene)camphor, 3-benzylidenecamphor;
- 4-aminobenzoic acid derivatives, preferably 2-ethylhexyl 4-(dimethylamino)benzoate, amyl 4-(dimethylamino)benzoate;
- esters of cinnamic acid, preferably 2-ethylhexyl 4-methoxycinnamate, isopentyl 4-methoxycinnamate;



- Examples of advantageous water-soluble UVB filter substances are:

- The list of said further UVB filters which can be used in combination with the active ingredient combinations according to the invention is of course not intended to be limiting.

It can also be advantageous to combine the combinations according to the invention with further UVA filters which have hitherto been customarily present in cosmetic preparations. These substances are preferably derivatives of dibenzoylmethane, in particular 1-(4'-tert-butylphenyl)-3-(4'-methoxyphenyl)propane-1,3-dione and 1-phenyl-3-(4'-isopropylphenyl)propane-1,3-dione. These combinations and preparations which contain these combinations are also provided by the invention. The amounts which may be used are as for the UVB combination.

It is further advantageous to combine the active ingredient combinations according to the invention with further UVA and/or UVB filters.

It is also particularly advantageous to combine the active ingredient combinations according to the invention with salicylic acid derivatives, some of which, as is known, can also absorb UV radiation. Customary UV filters include



The invention further provides a process for the preparation of the cosmetic and/or dermatological light protection preparations according to the invention, which comprises suspending and, if desired, homogenizing tris(2-ethylhexyl) 4,4',4''-(1,3,5-triazine-2,4,6-triyltriimino)trisbenzoate in one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols or an oil phase containing one or more esters of unbranched-chain carboxylic acids and branched-chain alcohols with uniform stirring and if necessary with warming, where appropriate combining the mixture with further lipid components and optionally with one or more emulsifiers, then mixing the oil phase with the aqueous phase into which a thickener has optionally been incorporated and which is preferably at the same temperature as the oil phase, if desired homogenizing the mixture, and allowing it to cool to room temperature. After the mixture has cooled to room temperature, homogenization may be repeated, particularly if volatile constituents are still to be incorporated.

The examples below serve to illustrate the present invention without limiting it. Unless indicated otherwise, all amounts, proportions and percentages are

based on the weight and the total amount or on the total weight of the preparations.

Example 1

O/W emulsion

% by weight

Stearic acid	3.50
Glycerol	3.00
Cetylstearyl alcohol	0.50
Dicaprylyl ether	8.00
Uvinul®T150	5.00
Hexyldecyl laurate	12.00
Sodium hydroxide (45% strength)	0.33
Carbomer	0.20
Preservative	q.s.
Perfume	q.s.
Water, demin.	ad 100.00

Example 2

W/O emulsion

% by weight

Arlacel®989	5.50
Butylene glycol	5.00
Hexyldecyl laurate	12.00
Uvinul®T150	5.00
Cetylstearyl isononanoate	6.00
Carbomer	0.20
Preservative	q.s.
Perfume	q.s.
Water, demin.	ad 100.00

006011 E0100260

Example 3

Hydrodispersion gel

	% by weight
Carbomer	0.50
Butylene glycol	5.00
Hexyldecyl laurate	10.00
Sodium hydroxide (45% strength)	0.35
Uvinul®T150	5.00
Hydroxypropylcellulose	0.60
Preservative	q.s.
Perfume	q.s.
Water, demin.	ad 100.00

006011-EDT00260